

Anterior Cruciate Ligament (ACL) Reconstruction Pre-Operative Primer

[Dan Bien PT, OCS, CSCS, CPSS](#)

ACL injuries can impact athletes and patients in many ways. Rehabilitation after ACL injuries and surgery is effective to address the many impairments following an ACL injury. It is critical that the athlete or patient and their care team understand what surgery and rehabilitation will entail to optimize the chances of a successful recovery.

Should I do pre-op rehab or exercises and will it help my recovery?

Yes, you should do some [pre-operative rehabilitation](#) because there is significant evidence of benefits to your post-operative outcome and the knee with pre-operative exercise.^{1-4,27} This pre-operative rehabilitation can be supervised by a physical therapist or unsupervised on your own. Pre-operative physical therapy may be limited due to insurance considerations in some instances. We have created [written](#) and [video](#) informational resources for basic exercises and steps which should allow you to address some of the impairments that commonly develop following an ACL injury if you are unable to undergo a formal supervised pre-operative rehabilitation or “prehab” program.

There is often significant swelling, and related pain or discomfort at the knee and affected limb due to tissue trauma from an ACL injury. Tissue injury, swelling, and discomfort may lead to knee stiffness, lower extremity weakness, altered proprioception or position sense at the joint, and alterations in movement and gait patterns. Extensive research shows that restoring full range of motion at the knee prior to surgery improves post-operative outcomes.⁴⁻⁵ Based on this evidence, some surgeons will not perform the ACL surgery until knee range of motion is fully restored, pending any other secondary injuries that may prevent this. Further research shows that improving lower extremity strength prior to surgery may also improve post-operative outcomes.^{2,6,7} The quadriceps, hamstring, calf, and hip musculature are most often impacted by an ACL injury and associated movement compensations. Completing a basic pre-operative rehabilitation program to address these impairments may improve the likelihood of a successful post-operative outcome and limit further pre-operative functional limitations.

When will I begin physical therapy and rehabilitation following surgery?

Typically, most patients will begin supervised physical therapy anywhere from 2-14 days following surgery depending on the preference of your surgeon. It is imperative that you follow the surgeon’s preference and begin physical therapy during this window to limit post-operative complications, and to set yourself up for a successful outcome.

What can I do to set myself up for success post-operatively?

- Swelling is expected post-operatively due to the inflammatory process, invasiveness of the surgical procedure, and trauma to the knee joint and tissues. However, excessive and prolonged swelling can lead to stiffness and range of motion deficits, muscular weakness and inhibition, and pain or discomfort.⁸⁻¹¹ It is advisable to manage the limb swelling through limb elevation, cryotherapy or icing, use of prescribed medications, intermittent exercise, and compression garments or wraps as directed by your surgeon and physical therapist. Be aware that periods of prolonged standing and/or walking may also contribute to excessive swelling in the early post-operative period, and you should plan to modify your activity accordingly.
- Pain control is another critical element to a successful rehabilitation and surgical outcome. Prescribed and over the counter medications may have some side effects, however they are often a useful tool to limit swelling and pain that may prevent completion of the home exercise program, restorative sleep, etc. that are essential to healing and recovery. Temporary use of these medications may allow faster and more comprehensive recovery when they are used as directed. There are also several online tools to help manage pain through evidence-based strategies as well as meditation, breathing, pain science education, and other tools. Ice has also been shown to improve pain control and decrease pain medication usage in the early post-operative period.¹⁸⁻²⁰
- Patients should strive for a well-balanced diet to ensure adequate nutrition including additional protein intake to build and repair tissues, allow wound/incision repair, and preserve immune function. This may take on added importance as strength training progresses to fully ensure muscle fiber repair following more intense exercises that may trigger muscle fiber breakdown from progressive overload for strengthening.
- Although sleep may be interrupted in the early stages of post-operative recovery due to discomfort from the surgery, brace application at the involved limb, and other factors, attempting to obtain adequate restorative sleep is vital to your body's recovery both physically and mentally. The use of ice just before bed, prescribed medications, and brace use while sleeping may limit pain that could negatively impact sleep quality.
- Additionally, there is some research evidence that applying ice to the affected knee for 20 minutes prior to exercise may allow improved forced production for greater exercise effectiveness during home exercise program and physical therapy sessions.²⁶

What type of psychological and mental health effects can I expect?

Rehabilitation following ACLR is long and physically and mentally challenging at times. Recovery after ACLR is rarely a steady linear progression, which can be frustrating. Following an ACL

injury, patients may experience many different emotions or reactions.¹³⁻¹⁶ Frustration due to functional loss and limitations and a loss in independence may commonly result. Due to the lengthy rehabilitation and slow progress at times, patients may struggle to maintain motivation and consistency with the rehabilitation program. It is important to share these kinds of feelings and challenges with your providers, who can help address these issues and work closely with you and an interdisciplinary care team to help you overcome these challenges.

Additionally, many athletes may experience a loss of social interaction with teammates and even loss of personal identity as an athlete, which can have negative psychological consequences.¹³⁻¹⁶ For example, depression rates in athletes following ACL injury has been estimated at up to 40%.¹³ Even non-athletes may experience a loss of connection to activities, identities, and social connection during this time that can feel isolating. For any patients, the prospect of a prolonged rehab may be daunting and discouraging at times and they may worry about their ability to return to “normal.” These and other psychological issues resulting from injury may impact mental health.

It is important to stay as connected as possible to activities you enjoy and social support networks, and to recognize and properly address any mental health issues following ACL injury. Your UOI interdisciplinary team can work collaboratively to support you, and we have a network of mental health professionals that we can refer you to if needed to optimize your post-op outcome and success.

What should I expect on my first physical therapy visit?

Your first physical therapy visit will involve the provider reviewing your history and details of your surgical procedure, getting baseline range of motion and strength data, beginning some treatment, developing a home exercise program based on the surgery details and demonstrated impairments or deficits, and providing some education to begin a successful rehab journey. You should wear loose fitting fitness clothing or bring shorts to allow your provider to properly examine and treat your knee.

How long will I be in a brace or on crutches following surgery?

Most patients will discontinue the brace and crutches 1-3 weeks after surgery. However, this is variable depending on the specific surgical procedures performed, your individual progress with rehabilitation and meeting certain accepted criteria or benchmarks such as the ability to perform an independent straight leg raise, pain, motor control and stability on the affected limb, and gait pattern or movement competency . Please note that certain related surgical procedures that involve meniscal repair, articular cartilage, or other anatomical structures may require a longer period of bracing or restricted weight bearing to allow adequate healing and to

limit stress. Your surgeon and therapist will review individual considerations for crutch and brace discharge with you.

When can I shower or bathe?

It is advisable to sponge bathe for the first few days following surgery. If you do not have weight bearing post-operative precautions and you're comfortable weight bearing at the surgical leg, you may shower with the wounds adequately covered to stay dry. Typically, saran wrap or plastic trash bags wrapped tightly at the leg to waterproof the knee and protect the incisions work best. You want to avoid getting the incisions wet which may increase the risk of infection, and you will not be permitted to submerge the incisions in a bath or pool until all surgical incisions are completely closed and fully healed.

What will physical therapy (PT) involve?

For PT appointments, please dress in comfortable clothing you can bend, move, and exercise comfortably in, and that will allow your provider to examine and/or treat your knee. PT will involve exercises to address knee range of motion and stiffness, lower extremity strength of the surgical and non-surgical leg, gait and movement training, and in many cases manual therapy and stretching to help restore normal mobility. Your PT provider may choose to implement other modalities such as neuromuscular electrical stimulation (NMES), blood flow restriction (BFR) training, or other equipment and techniques to expedite your recovery. As you improve and the program progresses, the program will shift towards strength training, movement retraining, and sport-specific training focus.

How often will my physical therapy sessions be?

This is variable according to insurance restrictions and financial considerations with copays, deductible responsibilities, etc. Your therapist will do their best to effectively allocate and utilize your available PT visits based on some of these considerations as well as your progress. For those whose benefits may be limited, we have developed video and exercise resources to help you continue to progress even when you are not able to attend physical therapy in-person.

Do I need to purchase additional equipment or materials?

We do not mandate or require that you purchase additional equipment. Adhering to the provided education suggestions and regular prescribed PT attendance is typically sufficient to allow successful rehabilitation. Some patients have reported added benefits from integration of a [portable neuromuscular electrical stimulation \(NMES\)](#) unit to facilitate quadriceps strength recovery.¹⁷

There is some additional evidence that use of electrical stimulation devices or transcutaneous electrical nerve stimulation may assist not only in pain control, but also with quadriceps muscle activation, especially when combined with ice for focal knee joint cooling.²⁵ Cryotherapy and [cryocompression](#) devices have been shown to assist with improved pain and swelling control¹⁸⁻²¹ Your insurance plan may fully or partially cover some of these items with a written order from your surgeon, but you should check with your insurer.

Some patients also report that a stretching/yoga strap or [Stretch Out Strap™](#) helpful following surgery when knee mobility and strength are limited, and certain positions are difficult to tolerate.

What should I avoid with the surgical leg post-operatively?

Post-op ACLR patients should avoid pivoting on the affected limb in a weight bearing position with the foot fixed on the ground such as when turning or during transfers. We do want patients to resume normal activities as soon as possible and permissible, including returning to fitness training and programming. However, you should consult your provider prior to resuming, because there may be activities that are not indicated and may be harmful in the early post-operative period.

How long will PT last?

This will again vary from patient to patient depending on multiple factors including: individual goals, your insurance coverage, and your rehabilitation progress.

Will I need to join a gym or perform weight training exercises?

Body weight exercises will be useful earlier in rehabilitation, but to fully restore strength at the affected leg, you will need to progressively overload the lower extremities with resistance exercises. Lower extremity weakness and lack of force production, specifically at the quadriceps is a common deficit following ACLR. For patients with a limited training history and limited familiarity with weight training technique, it may be useful to review technique with your physical therapist and/or implement [video resources](#). Additionally, it will be critical to work with your physical therapist to adjust the training volume and resistance to facilitate continued progress. Those patients with a limited training history may also need assistance to determine appropriate training intensity.

When can I return to work?

This will be a very individualized decision for each person based on job duties and requirements, financial considerations, job security and job stability concerns, and vacation or sick time available. Be aware that a return to a more physically demanding job may not be appropriate for several weeks or months following surgery. Also, jobs requiring prolonged

standing, walking, or positioning the leg in a dependent position for lengthy periods may contribute to an increase in swelling and discomfort, which may delay rehabilitation progress and recovery.

When can I begin to run again?

Running often begins between 3-5 months post-operatively. However, this will vary between individuals based on whether and when they attain established clinical benchmarks and criteria.

When can I return to sport?

Return to sports clearance is based on multiple considerations. However, time from surgery is not as important as meeting established clinical benchmarks and criteria. Benchmark testing may include: hop testing, vertical jump testing, peak force testing of specific muscle groups, torque/body weight ratio, movement/motor control testing, and other screening tools. 9 to 12 months is the typical timeframe to fully resume return to pivoting and/or contact sports, but this will vary between individuals.

Will I get back to my pre-surgery functioning?

A significant proportion of patients will resume pre-injury activities and performance levels.²²⁻²³ The timeframe for everyone to achieve these goals varies significantly and each patient's rehabilitation is different. It is often tempting to compare your progress to others who have previously undergone similar surgery. However, everyone's rehabilitation journey is unique and impacted by many factors. Working closely with your providers to identify, continually reassess, and target your specific deficits or impairments is critical to your success. Fully completing rehabilitation and achieving established clinical rehabilitation benchmarks and criteria significantly improves chances of return to sport and/or return to pre-injury activity and performance.

REFERENCES

1. Benefits of pre-op PT on ROM and outcome- particularly knee ext ROM. Loss of > 3 degrees impact mechanics.
Cunha J, Solomon DJ. ACL Prehabilitation Improves Postoperative Strength and Motion and Return to Sport in Athletes. *Arthrosc Sports Med Rehabil*. 2022 Jan 28;4(1):e65-e69. doi: 10.1016/j.asmr.2021.11.001. PMID: 35141537; PMCID: PMC8811524.
2. Riesterer J, Mauch M, Paul J, Gehring D, Ritzmann R, Wenning M. Relationship between pre- and post-operative isokinetic strength after ACL reconstruction using hamstring autograft. *BMC Sports Sci Med Rehabil*. 2020 Oct 30;12(1):68. doi: 10.1186/s13102-020-00215-7. PMID: 33292502; PMCID: PMC7602313.
3. <https://acltear.info/anterior-cruciate-ligament-rehabilitation/acl-prehab/#:~:text=Strengthen%20your%20muscles,your%20knee%20to%20promote%20healing.>
4. Quelard B, Sonnery-Cottet B, Zayni R, Ogassawara R, Prost T, Chambat P. Preoperative Factors Correlating with Prolonged Range of Motion Deficit after Anterior Cruciate Ligament Reconstruction. *The American Journal of Sports Medicine*. 2010;38(10):2034-2039. doi:10.1177/0363546510370198
5. Mauro CS, Irrgang JJ, Williams BA, Harner CD. Loss of extension following anterior cruciate ligament reconstruction: analysis of incidence and etiology using IKDC criteria. *Arthroscopy*. 2008 Feb;24(2):146-53. doi: 10.1016/j.arthro.2007.08.026. Epub 2007 Nov 5. PMID: 18237697.
6. Lepley LK, Palmieri-Smith RM. Pre-operative quadriceps activation is related to post-operative activation, not strength, in patients post-ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2016 Jan;24(1):236-46. doi: 10.1007/s00167-014-3371-0. Epub 2014 Oct 15. PMID: 25315083; PMCID: PMC4758128.
7. Logerstedt D, Lynch A, Axe MJ, Snyder-Mackler L. Pre-operative quadriceps strength predicts IKDC2000 scores 6 months after anterior cruciate ligament reconstruction. *Knee*. 2013 Jun;20(3):208-12. doi: 10.1016/j.knee.2012.07.011. Epub 2012 Sep 27. PMID: 23022031; PMCID: PMC3535501.
8. Andrä K, Kayaalp E, Prill R, Irlenbusch L, Liesaus E, Trommer T, Ullmann P, Becker R. Joint effusion, anteroposterior stability, muscle strength and degree of patellofemoral osteoarthritis significantly impact outcome following revision ACL reconstruction. *J Exp Orthop*. 2021 Aug 26;8(1):70. doi: 10.1186/s40634-021-00370-x. PMID: 34436684; PMCID: PMC8390615.

9. Kikuchi N, Kanamori A, Arai N, Okuno K, Yamazaki M. Joint Effusion at 3 Months After Anterior Cruciate Ligament Reconstruction: Its Risk Factors and Association With Subsequent Muscle Strength and Graft Remodeling. *Orthop J Sports Med.* 2024 Dec 16;12(12):23259671241299782. doi: 10.1177/23259671241299782. PMID: 39697608; PMCID: PMC11653272.
10. Rice DA, McNair PJ, Lewis GN, Dalbeth N. Quadriceps arthrogenic muscle inhibition: the effects of experimental knee joint effusion on motor cortex excitability. *Arthritis Res Ther.* 2014 Dec 10;16(6):502. doi: 10.1186/s13075-014-0502-4. PMID: 25497133; PMCID: PMC4271337.
11. Palmieri-Smith RM, Kreinbrink J, Ashton-Miller JA, Wojtys EM. Quadriceps inhibition induced by an experimental knee joint effusion affects knee joint mechanics during a single-legged drop landing. *Am J Sports Med.* 2007 Aug;35(8):1269-75. doi: 10.1177/0363546506296417. Epub 2007 Jan 23. PMID: 17244901.
12. Palmieri-Smith RM, Kreinbrink J, Ashton-Miller JA, Wojtys EM. Quadriceps inhibition induced by an experimental knee joint effusion affects knee joint mechanics during a single-legged drop landing. *Am J Sports Med.* 2007 Aug;35(8):1269-75. doi: 10.1177/0363546506296417. Epub 2007 Jan 23. PMID: 17244901.
13. Garcia GH, Wu HH, Park MJ, Tjoumakaris FP, Tucker BS, Kelly JD 4th, Sennett BJ. Depression Symptomatology and Anterior Cruciate Ligament Injury: Incidence and Effect on Functional Outcome--A Prospective Cohort Study. *Am J Sports Med.* 2016 Mar;44(3):572-9. doi: 10.1177/0363546515612466. Epub 2015 Dec 1. PMID: 26628516.
14. Burland JP, Toonstra JL, Howard JS. Psychosocial Barriers After Anterior Cruciate Ligament Reconstruction: A Clinical Review of Factors Influencing Postoperative Success. *Sports Health.* 2019 Nov/Dec;11(6):528-534. doi: 10.1177/1941738119869333. Epub 2019 Aug 30. PMID: 31469614; PMCID: PMC6822210.
15. Burland JP, Howard JS, Lepley AS, DiStefano LJ, Lepley LK, Frechette L. What Are Our Patients Really Telling Us? Psychological Constructs Associated With Patient-Reported Outcomes After Anterior Cruciate Ligament Reconstruction. *J Athl Train.* 2020 Jul 1;55(7):707-716. doi: 10.4085/1062-6050-120-19. PMID: 32702111; PMCID: PMC7384464.
16. Zuk EF, Kim S, Burland JP, Glaviano NR. The Comparison of Psychological Barriers Between Individuals with a History of Anterior Knee Pain, Anterior Cruciate Ligament Reconstruction, and Healthy Individuals. *Int J Sports Phys Ther.* 2023 Feb 1;18(1):92-101. doi: 10.26603/001c.68045. PMID: 36793558; PMCID: PMC9897036.
17. Arhos EK, Ito N, Hunter-Giordano A, Nolan TP, Snyder-Mackler L, Silbernagel KG. Who's Afraid of Electrical Stimulation? Let's Revisit the Application of NMES at the Knee. *J Orthop Sports Phys Ther.* 2024 Feb;54(2):101-106. doi: 10.2519/jospt.2023.12028. PMID: 37904496; PMCID: PMC10872626.

18. Waterman B, Walker JJ, Swaims C, Shortt M, Todd MS, Machen SM, Owens BD. The efficacy of combined cryotherapy and compression compared with cryotherapy alone following anterior cruciate ligament reconstruction. *J Knee Surg.* 2012 May;25(2):155-60. doi: 10.1055/s-0031-1299650. PMID: 22928433
19. Murgier J, Cassard X. Cryotherapy with dynamic intermittent compression for analgesia after anterior cruciate ligament reconstruction. Preliminary study. *Orthop Traumatol Surg Res.* 2014 May;100(3):309-12. doi: 10.1016/j.otsr.2013.12.019. Epub 2014 Mar 25. PMID: 24679367.
20. Yang JH, Hwang KT, Lee MK, Jo S, Cho E, Lee JK. Comparison of a Cryopneumatic Compression Device and Ice Packs for Cryotherapy Following Anterior Cruciate Ligament Reconstruction. *Clin Orthop Surg.* 2023 Apr;15(2):234-240. doi: 10.4055/cios21246. Epub 2022 Aug 26. PMID: 37008961; PMCID: PMC10060780.
21. Culvenor AG, Girdwood MA, Juhl CB, Patterson BE, Haberfield MJ, Holm PM, Bricca A, Whittaker JL, Roos EM, Crossley KM. Rehabilitation after anterior cruciate ligament and meniscal injuries: a best-evidence synthesis of systematic reviews for the OPTIKNEE consensus. *Br J Sports Med.* 2022 Dec;56(24):1445-1453. doi: 10.1136/bjsports-2022-105495. Epub 2022 Jun 29. PMID: 35768181; PMCID: PMC9726950.
22. Ardern CL. Anterior Cruciate Ligament Reconstruction-Not Exactly a One-Way Ticket Back to the Preinjury Level: A Review of Contextual Factors Affecting Return to Sport After Surgery. *Sports Health.* 2015 May;7(3):224-30. doi: 10.1177/1941738115578131. PMID: 26131299; PMCID: PMC4482306.
23. Ardern CL, Webster KE, Taylor NF, Feller JA. Return to sport following anterior cruciate ligament reconstruction surgery: a systematic review and meta-analysis of the state of play. *Br J Sports Med.* 2011 Jun;45(7):596-606. doi: 10.1136/bjism.2010.076364. Epub 2011 Mar 11. PMID: 21398310.
24. Kotsifaki R, King E, Bahr R, Whiteley R. Is 9 months the sweet spot for male athletes to return to sport after anterior cruciate ligament reconstruction? *Br J Sports Med.* 2025 Feb 26;bjsports-2024-108733. doi: 10.1136/bjsports-2024-108733. Epub ahead of print. PMID: 40011017.
25. Pietrosimone BG, Hart JM, Saliba SA, Hertel J, Ingersoll CD. Immediate effects of transcutaneous electrical nerve stimulation and focal knee joint cooling on quadriceps activation. *Med Sci Sports Exerc.* 2009 Jun;41(6):1175-81. doi: 10.1249/MSS.0b013e3181982557. PMID: 19461552.
26. Hart JM, Kuenze CM, Diduch DR, Ingersoll CD. Quadriceps muscle function after rehabilitation with cryotherapy in patients with anterior cruciate ligament reconstruction. *J Athl Train.* 2014 Nov-Dec;49(6):733-9. doi: 10.4085/1062-6050-49.3.39. PMID: 25299442; PMCID: PMC4264644.

27. Failla MJ, Logerstedt DS, Grindem H, Axe MJ, Risberg MA, Engebretsen L, Huston LJ, Spindler KP, Snyder-Mackler L. Does Extended Preoperative Rehabilitation Influence Outcomes 2 Years After ACL Reconstruction? A Comparative Effectiveness Study Between the MOON and Delaware-Oslo ACL Cohorts. *Am J Sports Med.* 2016 Oct;44(10):2608-2614. doi: 10.1177/0363546516652594. Epub 2016 Jul 14. Erratum in: *Am J Sports Med.* 2017 Apr;45(5):NP9. doi: 10.1177/0363546517696318. PMID: 27416993; PMCID: PMC5537599.